



1100 East Sandy Lake Road  
Coppell, TX 75019-3113  
972.471.7775

---

## **TOMATOES, CUCUMBERS, PEPPERS** **(field and greenhouse)**

### **General Application Rates/Methods:**

#### ***Direct seed planting:***

***Soil:*** Sandy soil 1 quart to the acre, Clay or high organic matter soil 12.8 ounces to the acre. Sandy loam 1 gallon when plants are in full flowering

- In furrow with seed at planting if possible

***Foliar (drench):*** 1 quart to the acre at 4-6 leaf

#### ***Transplants:***

***Foliar (drench):*** 12.8 oz to the acre  
Repeat at start of bloom (flowering)

***Soil:*** 1 quart to the acre at fruit set

### **TOMATOES (indeterminate)**

***Planting:*** Soil application: Sandy soil 1 quart to the acre, Clay or high organic matter soil 12.8 ounces to the acre. Sandy loam 1 gallon when plants are in full flowering.

Repeat soil application every two weeks until flowering.

***Foliar:*** Once per month after third inflorescences

### **TOMATOES (determinate)**

See General Application above.

## **TOMATOES, CUCUMBERS, PEPPERS**

### **Page 2**

#### **Benefits:**

Improved storability.

Improved food value – high in nutrients and lycopene

Better brix -- improved acid to sugar ratios

Improved vine ripening

#### **For Best Results in Agricultural Application:**

**Applied thru overhead, aerial irrigation and mixed with sprayer tank containing start up fertilizers and/or herbicides.**

Ensure tank is free of residue, avoid chlorinated water if possible.

Do not apply when a crop is severely stressed.

Do not mix with fungicides or insecticides and do not apply within 2 weeks of applying fungicides.

May be mixed with a 2,4-D herbicide, and Round-up as well as roundup ready seeds.

In light of new seed developments and/or stacked seeds, contact manufacturer's technical support.